



Exosomal therapy – a new frontier in regenerative medicine

Sathish Muthu^{1,2,3^}, Asawari Bapat^{4^}, Rashmi Jain^{3,5^}, Naveen Jeyaraman^{3,6^}, Madhan Jeyaraman^{2,3,7^}

¹Department of Orthopaedics, Government Hospital, Velayuthampalayam, Karur, Tamil Nadu, India; ²Department of Biotechnology, School of Engineering and Technology, Sharda University, Greater Noida, Uttar Pradesh, India; ³Indian Stem Cell Study Group (ISCSG), Lucknow, Uttar Pradesh, India; ⁴Director of Quality and Regulatory Affairs, Infohealth FZE, Dubai, United Arab Emirates; ⁵School of Medical Sciences and Research, Sharda University, Greater Noida, Uttar Pradesh, India; ⁶Department of Orthopaedics, Kasturba Medical College, MAHE University, Manipal, Karnataka, India; ⁷Department of Orthopaedics, School of Medical Sciences and Research, Sharda University, Greater Noida, Uttar Pradesh, India

Contributions: (I) Conception and design: A Bapat; (II) Administrative support: N Jeyaraman; (III) Provision of study materials or patients: M Jeyaraman, S Muthu; (IV) Collection and assembly of data: M Jeyaraman, S Muthu; (V) Data analysis and interpretation: R Jain; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Dr. Madhan Jeyaraman. Department of Orthopaedics, School of Medical Sciences and Research, Sharda University, Greater Noida, Uttar Pradesh, India. Email: madhanjeyaraman@gmail.com.

Abstract: The recent advances in translational and nanomedicine have paved the way for developing the targeted drug delivery system at a greater pace among global researchers. On par with these technologies, exosomes act as a potential portal for cell-free drug delivery systems as these are bestowed with the native characteristics of the parent cell of origin. Exosomes, called extracellular vesicles (EcVs), are present in almost all cells, tissues, and body fluids. They help in intercellular signaling and maintains tissue homeostasis in the disease pathobiology. Researchers have characterized 9,769 proteins, 2,838 miRNAs, 3,408 mRNAs, and 1,116 lipids being present in exosomal cargo. The separation of exosomes from cells, tissues, and body fluids follow different patterned kinetics. Exosomes interact with the recipient cells through their surface receptor molecules and ligands and internalize within recipient cells through micropinocytosis and phagocytosis. Advancing technologies in regenerative medicine have facilitated the researchers to isolate exosomes from mesenchymal stem cells (MSCs) as these cells are blessed with supreme regenerative potentiality in targeting a disease. Exosomal cargo is a key player in establishing the diagnosis and executing therapeutic role whilst regulating a disease process. Various *in vitro* studies have exhibited the safety, efficacy, and therapeutic potentiality of exosomes in various cancers, neurodegenerative, cardiovascular, and orthopedic diseases. This article throws light on the composition, therapeutic role, and regulatory potentials of exosomes with the widening of the horizon in the field of regenerative medicine.

Keywords: Exosomes; extracellular vesicles (EcVs); cellular therapy; biological medicine

Received: 03 August 2020; Accepted: 16 March 2021; Published: 02 April 2021.

doi: 10.21037/sci-2020-037

View this article at: <http://dx.doi.org/10.21037/sci-2020-037>

Introduction

The contemporary developments in the field of translational medicine include the development of targeted drug delivery systems for harnessing the full potentiality and

utilizing the therapeutic products effectively by overcoming the limitations of the existing methods to address the pathogenesis of a disease. In that context, various polymer-based nano delivery systems have been developed (1). Utilizing extracellular vesicles (EcVs) as a drug delivery

[^] ORCID: Sathish Muthu, 0000-0002-7143-4354; Asawari Bapat, 0000-0002-4592-7702; Rashmi Jain, 0000-0003-4386-6755; Naveen Jeyaraman, 0000-0002-4362-3326; Madhan Jeyaraman, 0000-0002-9045-9493.

tool has been given special attention due to their naïve characteristics derived from the parent or host cells (2-4). Despite the normal cellular homeostasis, EcVs play a major role in intervening in the pathobiology of disease processes through the intercellular signaling cascade (5-9).

Based on the size, exosomes are subcategorized under EcVs, which are endosome derived lipid bi-layered spherical vesicles of 40–150 nm in size (2,10-12). Almost all cells, tissues, and body fluids [plasma, urine, saliva, tears, gastrointestinal (GI) secretions, semen, and breast milk] secrete exosomes (13-16). Exosomal cargo carries an array of micro-biomolecules that consists of proteins, lipids, RNA, and DNA from the secreting parent cells (2,17-19). Moreover, the characteristics and behavior of the exosomes closely relate to the parent cell of origin (20-22). On account of their appropriate size and property along with their evident role in numerous pathobiological processes, the potential of exosomal therapy in the management of various neurodegenerative disorders, infectious diseases, musculoskeletal disorders, and cardiovascular disorders is overwhelming. This has enabled the researchers to target the disease process at a cellular level by using natural engineered defense mechanisms.

Characterization of exosomes

The composition of an exosome vesicle (EV) includes proteins, RNA, DNA, and other substances. Currently, 9,769 proteins, 2,838 miRNAs, 3,408 mRNAs, and 1,116 lipids have been described in their composition (23). Of these components, the exosomal proteins differ based on the nature of the primitive cells or tissues (24). Proteins like membrane transport and fusion proteins, chaperones, adhesion molecules, MHCs, cytoskeletal proteins, and lipid-related proteins are major exosomal proteins (24-28). With a pre-existing knowledge of the functioning of these proteins, ALG-2-interacting protein X (ALIX), heat shock protein 70 (HSP 70), tumor susceptibility gene 101 (TSG101), and tetraspanins (CD9, CD 63, CD 81, and CD 82) have been identified with a higher concentration in exosomes (29,30). Moreover, they serve as a biomarker for exosomal identification as shown in *Figure 1* (31). Exosomes are also rich in lipid layering molecules like glycosylphosphatidylinositol-anchored protein (LBPA) and flotillin (32,33). Apart from the metabolic enzymes and signal transduction molecules such as G protein and protein kinases, exosomes contain mRNAs, miRNAs, non-coding RNAs (ncRNAs), and mitochondrial DNA in

their composition (34,35). Noteworthy, the first types of nucleic acids identified in the exosomes were mRNA and miRNA (36,37).

Being a nanoscale vesicular component in complex body fluids, it is challenging to procure exosomes of high-yield and fineness (38). However, their isolation and characterization is an essential prerequisite to establish their therapeutic potential with a better understanding of their physiology. Exosomes are derived from the cell culture supernatants or plasma with their identification based on the physical and morphological characteristics (39,40). Although separation can be made by several methods like ultracentrifugation, ultrafiltration, gradient ultracentrifugation, precipitation, size-exclusion chromatography, immune-affinity capture, mass spectrometric immunoassay, magnetic-activated cell sorting, and microfluidics-based techniques with each having their advantages and disadvantages (41-45) Ultracentrifugation is the most commonly used method for isolating exosomes. The complex proteins in the exosomes are analyzed by processes like western blot, flow cytometry, and mass spectrometry (46). Additionally, sample analysis with high precision fluorescence is a possibility in the nanoparticle tracking method (47). Homogeneity without exosome isolation is possible with the resistance pulse sensing method (48). Profiling the miRNA content of the exosomes was made with next-generation sequencing, microarray processing, and RT-PCR. Confirmation of the isolated samples can be done by electron microscopy (49). Disparate use of various methods of isolation, confirmation, quantification, and analysis used by the researchers with their own set of advantages and disadvantages, it brings added heterogeneity in their analysis which needs standardization.

Cellular physiology of exosomes

The formation of exosomes involves the invagination of the plasma membrane and the formation of intracellular multivesicular bodies with intraluminal vesicles. This endocytic pathway of the donor cell is followed by the transport of the transmembrane and intra-vesicular proteins from the Golgi complex resulting in the formation of early endosomes. After maturation and differentiation, they get transformed into late endosomes (50-53). They are degraded by fusing with lysosomes or plasma membrane or autophagosomes to release the intraluminal vesicles as exosomes (measuring 40–150 nm diameter) into the extracellular milieu as shown in *Figure 2* (5,32,54).




Figure 1 Basic structure of mesenchymal stem cell exosome depicting the surface and core proteins along with the key immune and growth factors.

Exosomes interact with the recipient cells through their surface receptor molecules and ligands. Some of the exosomes persist on the cell membranes of the donor cells after secretion while the rest of them interacts with the recipient cells (54-59). Internalization of exosomes occurs via membrane integration process mediated by raft or caveolae or clathrin dependent endocytosis. Micropinocytosis and phagocytosis have also been described as a method of internalization of the exosomes by recipient cells. This physiological integration process on the targeted recipient cells has been considered to have therapeutic potential as a targeted delivery system for effectively executing biological functions (60-64). Yet, the exosomal components in specific accounting for cell-type or organ specificity remain unclear (65).

Diagnostic role of exosomes

Exosomes are the key players of the intercellular communication portal and thereby owe the capability of determining the progression of a disease. Studies are emerging in the diagnostic and therapeutic facets of exosomes for various systemic pathologies. The detection of substances (intracellular and extracellular) carried by these nanoparticles and promoting their immune capture with surface proteins aid in the diagnosis of pathological processes. The spectrum of diseases in which exosomes play a key role in diagnosis includes cerebrovascular disease (66,67), diseases involving the central nervous system and neoplasm (8,68-70) along with disease involving kidney, liver, and lungs (70-72). Let us consider the scenario of cancer. Exosomes can protect the rapid degradation of




Figure 2 Physiology of formation and degradation of exosomes. ECM, extracellular matrix.

nucleic acids and remain highly stable in plasma (73). With an exosome-based liquid biopsy, direct detection of the circulating tumor cells and their DNA or cell-free RNA from body fluids like blood, saliva can be made. This DNA or RNA (including mRNA, miRNA, ncRNA) from serum exosomes can be used for the detection of cancer-related mutations making them a promising biomarker in cancer diagnosis (74-77). Various studies have identified proteins and RNAs rich in cancer-derived exosomes which act as biomarkers that can be used in the detection and prognosticating treatment response (78-85).

Therapeutic role of exosomes

Exosomes possess a higher therapeutic potential for various disease spectra due to their ability for intracellular shuttling. Nanomedicine technologies have given rise to explore

the usage of pathogenic importance of exosomal particles in various diseases. The targeted drug delivery system in nanomedicine focus on the sustained release of exosomes for exerting the biological activity in the targeted site. Exosomes are used as vectors or carrier molecules to elicit a biological response.

Under given physiological circumstances, exosomes demonstrate very low immunogenicity and the potential to circumvent the physiological blood-brain-barrier (86). With the help of a stable lipid bilayer, the cargoes loaded in the exosomal vesicles are guarded against the action of native immune cells and digestive enzymes. The engineered exosomal vesicles deliver the cargoes loaded to them to the site of action through various mechanisms of endocytosis or membrane fusion as shown in *Figures 3 and 4* (87,88). EVs are derived from various cell types and tissues. On delivery to specific diseased tissue, under specific conditions, EV elicits




Figure 3 Methods of uptake of exosomes by the target cells.




Figure 4 Potential therapeutic applications of exosomes. ILD, interstitial lung disease; IBD, inflammatory bowel disease.

tissue regeneration and homeostasis (89). Mesenchymal stromal cell-derived EV show cellular viability, tropicity, anti-inflammatory, immunomodulatory and therapeutic effects (90,91). They support neoangiogenesis and cellular proliferation (2,92-95). Exosomes demonstrate the homing effect of the parental cell type (2,96).

The superiority of the exosomal cargoes compared to the stem cells on clinical and therapeutic potential is noted by

- (I) Lack of inherent risk associated with any cell-based therapies including stem cells;
- (II) Lack of replicating potential and the risk of malignant transformation;
- (III) Lack of immunogenic response towards infections and cancers;
- (IV) Targeted action at the site of interest (2,97).

The above-mentioned versatility of EVs enhances their intracellular signaling capabilities and their movement across cellular membranes to restore micromolecular homeostasis. Apart from these advantages, exosomes also offer neuroprotection and neuroplasticity by shuttling across the blood-brain barrier in neurodegenerative diseases (98). A total of 77 clinical trials have been registered among the global researchers in various pathologies. Out of 77 trials, 11 clinical trials have been completed and have proved the positive diagnostic and therapeutic effects of exosome-based treatment in various cancers, neurodegenerative and hematological diseases (99).

Sources of isolation of exosomes

The exosomes are also called intraluminal vesicles and are found in various tissues and body fluids (5-7). The exosomes derived from mesenchymal stem cells (MSCs) are of prime importance due to the greater therapeutic and regenerative potential as shown in *Figure 4*. Due to the challenges faced in isolating exosomes from various body fluids, regenerative and translational medicine experts used MSC-derived exosomes for treating various disorders (2). MSCs derived exosomal cargo exhibit intracellular signaling and communications to targeted tissues. The key sources of exosomes derived from the MSCs include bone marrow, adipose tissue, placental cells, umbilical cells, endometrial fluid, and amniotic fluid (2,100).

Exosomes of MSC origin have cell surface markers such as CD 29, CD 44, and CD 73 embedded in them (100). They play a vital role in the biomechanisms involved in the repair and regeneration, bioenergetics, immunoregulation, intracellular communication, and tissue metabolism (2,101).

In a proteomic analysis, a total of 730 protein molecules in bone marrow MSC derived exosomes were isolated (102). Some researchers found the existence of transcription signaling factors in exosomal cargoes (103). Amniotic fluid exosomes are the most preferred exosomes for clinical applications than bone marrow-derived exosomes (104).

Routes of delivery of exosomes

The different state of the art approaches to deliver the exosomes to their site of action and their shortcomings were analyzed and presented with their challenges encountered in them. The most common route is intravenous (IV) despite its clearance in the liver and the kidney is rapid in this method (105). This method was widely used for conditions involving orthopedic, neoplastic, and cardiovascular pathologies (106). Intramuscular (IM) has been primarily used in neuromuscular and musculoskeletal conditions while subcutaneous (SC) route in aesthetic and cosmetic indications. The IM or SC route is chosen due to the ease of injectable area and the dosing volume (97). Intrathecal is the preferred route in neurodegenerative conditions such as Alzheimer's, Parkinson's, and Creutz-Feld Jakob's disease (98,99). Local aerosol sprays are used in the management of wounds and ulcers (107,108). For hair growth and rejuvenation in age-related therapies, this is the preferred route (108,109). During the time of the coronavirus disease-2019 (COVID-19) pandemic, there have been interesting studies discussing the role of exosomal therapy as a promising therapeutic candidate (110).

Global regulatory requirements

With the growing therapeutic spectrum of exosomal therapy, the International Society for Extracellular Vesicles (ISEV) and the European Network on Microvesicles and Exosomes in Health and Disease (ME-HaD) have formulated certain guidelines to foster their clinical usage (10). The regulations elaborate on the standard operative protocols to be followed in the process of collection, processing, testing, quality control, and manufacturing of exosomes for clinical usage. With the help of these policies, the potential of the EVs can be utilized at appropriate standards for therapeutic usage.

Any novel therapy or a drug being developed is dependent on the strategies to standardize the process to focus on validating the proposed technology. There are currently no Food and Drug Administration (FDA)-

approved exosome products for human use in the USA (2,111). According to the FDA, exosomes are classified as a 351 product that requires studies effectively showing safety and efficacy, along with the purity of the product and its potency in treating the condition (112). Therapies using the exosomes are under the Investigational New Drug (IND) developmental phase and need the approval of the regulatory agencies before initiating the clinical trial (2,113).

The regulatory framework addresses the safety standards for microbial and viral contamination and demands GxP standards (GxP = good manufacturing/good laboratory/good distribution/good clinical/good scientific practice or GMP/GLP/GDP/GCP/GSP) for the production and quality control of the corresponding therapeutics (114,115). It regulates the conduct of clinical trials (116). There remains an issue with the categorization of any emerging novel therapeutics for humans.

Pharmaceutical category of EV preparations

According to the Center for Biologics Evaluation and Research (CBER), the exosomes are regulated as biological products (117,118). Based on the individual types, the framework that was laid down for products in this category applies to the exosomes. For example, an anti-tumor vaccine that uses exosomes will be regulated under the regulations provided for therapeutic cancer vaccines (119). The functional moiety in EV-based therapy determines its medicinal type (2,120).

Hence, ISEV categorizes EV-based therapy under biological medicines with the following properties.

- (I) The therapeutics acquired from unmodified cells.
- (II) Therapeutics acquired from genetically manipulated cells (without trans-gene).
- (III) Therapies acquired from exosomes and gene-modified cells with trans-gene classified as gene therapy products (GTP).
- (IV) Native exosomal therapies; are used as drug-delivery systems, used as carriers for the biological and chemical components, and are considered as biological medicine.

As the biological medicinal products include a span of various pharmaceuticals, these were classified as Advanced Therapy Medicinal Products (ATMP's) in 2007 (2,120). It was further subgrouped to conventional biological medicinal products due to the biological properties, physicochemical and immunochemical properties (117,118). It includes somatic cell therapy, gene therapies, and tissue-

engineered products.

ATMP's therapeutics involve products that have been:

- (I) More than minimally manipulated such as cell expansions and cell cultures.
- (II) Intended for non-homologous use. For example use of hematopoietic cells for orthopedics.
- (III) Nucleated and viable cells are present in the product.
- (IV) Products with therapeutically active trans-gene from genetically engineered cells are considered as ATMPs, independent of the presence of any nucleated, viable cell.
- (V) EV-based therapeutics classified as ATMP's; are produced from human material by a manufacturing process comparable to the ATMP production.

Safety profile, manufacturing & standardization

The mechanism of action (MoA) is essential for the clinical translation of therapeutics based on EVs (121). The critical part of this translation is the identification of "active substances", their properties, and the essential quality controls in manufacturing a clinical-grade product (118). For phase I the hypothesis should be based on the proof of principle, reinstate the rationale based on the MoAs. Although uncertainties exist on the MoA of EV-based therapeutics on the target cells, with supportive animal models, EVs from human cells are not to be accounted as high risk in an IND (116). Commercializing a large scale manufacturing of EV based therapeutics requires a robust quality management system, technologically superior facility, and an updated technology complying with GxP (114,115). The endpoint of these investigational studies is to provide safety for the donor and the patient. The criteria for the therapeutic release of an investigational product is to determine efficacy based on the pre-IND studies for characterization (116).

At this moment the EV's do not have a standardized protocol for isolation and storage; and include homemade cocktails as protocols with no standardization for reagents, storage containers, and storage time for each desired EV-based product (2,10,122).

Direction for future research

The scope of exosomal therapy among various other clinical fields remains untouched. With the global interest of the researchers towards harnessing the potential of

Table 1 Companies targeting on exosomal research and their potential products for commercial use

Company	Exosomal source	Exosomal product	Exosomal potential
Capricor Therapeutics (128)	Cardiosphere-derived cells (CDCs)	CAP-2003 (preclinical testing)	Explore the anti-inflammatory, pro-angiogenic, antiapoptotic and antifibrotic effects associated with their parent CDCs
Kimera Labs (129)	Placental MSC-derived exosome	XoGlo	Tap into the potential for tissue repositioning, preventing healing-associated scarring
Exosomedx (130)	Plasma or serum-derived exosomes	ExoDx	CLIA certified product to conduct advanced clinical testing
NanoSomiX (131)	Brain-derived exosomes (BDEs)	ExoM and ExoC	Biomarkers for prediction of potential nervous system disorders
Evox Therapeutics (132)			For rare metabolic and lysosomal storage diseases engaging endogenous and exogenous drug loading
Codiak Biosciences (133)	Engineered therapeutic exosomes		Encompasses therapeutics for vaccines, anti-infectives, oncology, autoimmune and anti-inflammatory diseases
Aruna Biomedical (134)	Neural derived exosomes	AB126	Murine thromboembolic models of stroke and its translation to clinical therapies

MSC, mesenchymal stem cell; CLIA, Clinical Laboratory Improvement Amendments.

the exosomes, making EV based therapeutics a reality is not far from reach (10). The potential of EV based therapy is established in various orthopedic conditions, neurodegenerative disorders, auto-immune diseases, cardiovascular diseases, infectious diseases, and diagnosis of rare diseases including cancers (98,99,123). However, the current domain of research in EV based therapeutics involve developing diagnostic and therapeutic applications towards patient care.

From the level of the circulating pool of EVs, early diagnosis of a complex disease could be made by using them as biomarkers in blood (74-82). Utilising the paracrine signaling property of the MSC-derived exosomes, repair, and regeneration of organs could be achieved (87-95). For use in most orthopedic conditions exosomes are deemed anti-inflammatory concerning their immunomodulatory potential (123). Understanding the role of exosomes in the disease processes have enabled us to propel our understanding of the disease and expand the scope of the therapies evolved out of them (124). As of now therapeutic applications of the EV-based therapies from defined cell sources based on their immunomodulatory capabilities include inflammatory disorders, degenerative disorders, vehicles of drug-delivery, anti-tumor therapies, and pathogen vaccination tool (98-102). Immunomodulation of exosomes may be exerted by either immune-activation or suppression. This novel platform with the above-mentioned diverse potentials holds promise to develop vaccines with

prolonged immunogenicity against infectious diseases or cancer (125,126).

There is a paradigm shift brought down by the continual breakthroughs in research exploring the potential of the exosomes that have resulted in the development of novel therapeutic options that are reshaping the landscape of the global market from time to time. Despite being started as a subject of academic interest, exosomal therapy has now been transformed into a potential platform with immense promise for future therapeutics (127). This sowed the seeds for start-ups into exosome platforms with their proprietary technologies. Some of them are elaborated in *Table 1*. These companies are using patented technologies to tap the potential of exosomes. For example, the company Capricor Therapeutics developed an exosomal therapeutic technology from cardiosphere-derived cells (CDCs) called CAP-2003 which is in the preclinical testing phase to explore the anti-inflammatory, pro-angiogenic, antiapoptotic, and antifibrotic effects associated with their parent CDCs. Additionally, the company is evaluating the role of CDC-exosomes for the treatment of trauma-related injuries and conditions (135).

Conclusions

Exosomes play a natural role by enacting as a vehicle for the transfer of biological substances between cells and thereby renders a broader prospect to serve as a channel

for delivering drugs of therapeutic interest. However, the intricate composition and uncertain functioning are inquisitive facets warranting further exploration. In view of making the exosome-based therapy a reality, more accurate, faster, cheaper, standardized, specific, and easier methods of their separation and purification have to evolve along with concrete adducing on its safety, feasibility, pharmacokinetic and pharmacodynamic characteristics through large scale prospective research studies.

Acknowledgments

Funding: None.

Footnote

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/sci-2020-037>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Wu D, Abezgauz L, Danino D, et al. Alternating polymer vesicles. *Soft Matter* 2008;4:1066-71.
2. Jeyaraman M, Muthu S, Gulati A, et al. Mesenchymal Stem Cell-Derived Exosomes: A Potential Therapeutic Avenue in Knee Osteoarthritis. *Cartilage* 2020. doi: 10.1177/1947603520962567.
3. Pata V, Dan N. The effect of chain length on protein solubilization in polymer-based vesicles (polymersomes). *Biophys J* 2003;85:2111-8.
4. Zhang XY, Zhang PY. Polymersomes in Nanomedicine - A Review. *Current Nanoscience* 2016. doi:10.2174/1573413712666161018144519.
5. Kalluri R, LeBleu VS. The biology, function, and biomedical applications of exosomes. *Science* 2020;367:eaau6977.
6. Pegtel DM, Gould SJ. Exosomes. *Annu Rev Biochem* 2019;88:487-514.
7. Becker A, Thakur BK, Weiss JM, et al. Extracellular Vesicles in Cancer: Cell-to-Cell Mediators of Metastasis. *Cancer Cell* 2016;30:836-48.
8. Kalluri R. The biology and function of exosomes in cancer. *J Clin Invest* 2016;126:1208-15.
9. Robbins PD, Morelli AE. Regulation of immune responses by extracellular vesicles. *Nat Rev Immunol* 2014;14:195-208.
10. Théry C, Witwer KW, Aikawa E, et al. Minimal information for studies of extracellular vesicles 2018 (MISEV2018): a position statement of the International Society for Extracellular Vesicles and update of the MISEV2014 guidelines. *J Extracell Vesicles* 2018;7:1535750.
11. Cocucci E, Meldolesi J. Ectosomes and exosomes: shedding the confusion between extracellular vesicles. *Trends Cell Biol* 2015;25:364-72.
12. O'Brien K, Breyne K, Ughetto S, et al. RNA delivery by extracellular vesicles in mammalian cells and its applications. *Nat Rev Mol Cell Biol* 2020;21:585-606.
13. Caby MP, Lankar D, Vincendeau-Scherrer C, et al. Exosomal-like vesicles are present in human blood plasma. *Int Immunol* 2005;17:879-87.
14. Pisitkun T, Shen RF, Knepper MA. Identification and proteomic profiling of exosomes in human urine. *Proc Natl Acad Sci U S A* 2004;101:13368-73.
15. Michael A, Bajracharya SD, Yuen PS, et al. Exosomes from human saliva as a source of microRNA biomarkers. *Oral Dis* 2010;16:34-8.
16. Admyre C, Johansson SM, Qazi KR, et al. Exosomes with immune modulatory features are present in human breast milk. *J Immunol* 2007;179:1969-78.
17. Mathivanan S, Ji H, Simpson RJ. Exosomes: extracellular organelles important in intercellular communication. *J Proteomics* 2010;73:1907-20.
18. Sokolova V, Ludwig AK, Hornung S, et al. Characterisation of exosomes derived from human cells by nanoparticle tracking analysis and scanning electron microscopy. *Colloids Surf B Biointerfaces* 2011;87:146-50.
19. Kalra H, Adda CG, Liem M, et al. Comparative proteomics evaluation of plasma exosome isolation techniques and

- assessment of the stability of exosomes in normal human blood plasma. *Proteomics* 2013;13:3354-64.
20. Théry C, Ostrowski M, Segura E. Membrane vesicles as conveyors of immune responses. *Nat Rev Immunol* 2009;9:581-93.
 21. Daaboul GG, Gagni P, Benussi L, et al. Digital Detection of Exosomes by Interferometric Imaging. *Sci Rep* 2016;6:37246.
 22. Pitt JM, Kroemer G, Zitvogel L. Extracellular vesicles: masters of intercellular communication and potential clinical interventions. *J Clin Invest* 2016;126:1139-43.
 23. Mathivanan S, Fahner CJ, Reid GE, et al. ExoCarta 2012: database of exosomal proteins, RNA and lipids. *Nucleic Acids Res* 2012;40:D1241-D1244.
 24. Mathivanan S, Lim JW, Tauro BJ, et al. Proteomics analysis of A33 immunoaffinity-purified exosomes released from the human colon tumor cell line LIM1215 reveals a tissue-specific protein signature. *Mol Cell Proteomics* 2010;9:197-208.
 25. Conde-Vancells J, Rodriguez-Suarez E, Embade N, et al. Characterization and comprehensive proteome profiling of exosomes secreted by hepatocytes. *J Proteome Res* 2008;7:5157-66.
 26. Poliakov A, Spilman M, Dokland T, et al. Structural heterogeneity and protein composition of exosome-like vesicles (prostasomes) in human semen. *Prostate* 2009;69:159-67.
 27. Simons M, Raposo G. Exosomes--vesicular carriers for intercellular communication. *Curr Opin Cell Biol* 2009;21:575-81.
 28. Subra C, Grand D, Laulagnier K, et al. Exosomes account for vesicle-mediated transcellular transport of activatable phospholipases and prostaglandins. *J Lipid Res* 2010;51:2105-20.
 29. Vlassov AV, Magdaleno S, Setterquist R, et al. Exosomes: current knowledge of their composition, biological functions, and diagnostic and therapeutic potentials. *Biochim Biophys Acta* 2012;1820:940-8.
 30. Xie F, Zhou X, Fang M, et al. Extracellular Vesicles in Cancer Immune Microenvironment and Cancer Immunotherapy. *Adv Sci (Weinh)* 2019;6:1901779.
 31. Jeppesen DK, Fenix AM, Franklin JL, et al. Reassessment of Exosome Composition. *Cell* 2019;177:428-445.e18.
 32. van Niel G, D'Angelo G, Raposo G. Shedding light on the cell biology of extracellular vesicles. *Nat Rev Mol Cell Biol* 2018;19:213-28.
 33. Valadi H, Ekström K, Bossios A, et al. Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nat Cell Biol* 2007;9:654-9.
 34. Guescini M, Genedani S, Stocchi V, et al. Astrocytes and Glioblastoma cells release exosomes carrying mtDNA. *J Neural Transm (Vienna)* 2010;117:1-4.
 35. Soung YH, Ford S, Zhang V, et al. Exosomes in Cancer Diagnostics. *Cancers (Basel)* 2017;9:8.
 36. Ratajczak J, Miekus K, Kucia M, et al. Embryonic stem cell-derived microvesicles reprogram hematopoietic progenitors: evidence for horizontal transfer of mRNA and protein delivery. *Leukemia* 2006;20:847-56.
 37. Turchinovich A, Weiz L, Langheinz A, et al. Characterization of extracellular circulating microRNA. *Nucleic Acids Res* 2011;39:7223-33.
 38. Willms E, Cabañas C, Mäger I, et al. Extracellular Vesicle Heterogeneity: Subpopulations, Isolation Techniques, and Diverse Functions in Cancer Progression. *Front Immunol* 2018;9:738.
 39. Bard MP, Hegmans JP, Hemmes A, et al. Proteomic analysis of exosomes isolated from human malignant pleural effusions. *Am J Respir Cell Mol Biol* 2004;31:114-21.
 40. Navabi H, Croston D, Hobot J, et al. Preparation of human ovarian cancer ascites-derived exosomes for a clinical trial. *Blood Cells Mol Dis* 2005;35:149-52.
 41. Li P, Kaslan M, Lee SH, et al. Progress in Exosome Isolation Techniques. *Theranostics* 2017;7:789-804.
 42. Momen-Heravi F, Balaj L, Alian S, et al. Current methods for the isolation of extracellular vesicles. *Biol Chem* 2013;394:1253-62.
 43. Théry C, Amigorena S, Raposo G, et al. Isolation and characterization of exosomes from cell culture supernatants and biological fluids. *Curr Protoc Cell Biol* 2006;Chapter 3:Unit 3.22. doi: 10.1002/0471143030.cb0322s30.
 44. Xu R, Greening DW, Zhu HJ, et al. Extracellular vesicle isolation and characterization: toward clinical application. *J Clin Invest* 2016;126:1152-62.
 45. Gulei D, Irimie AI, Cojocoreanu-Petric R, et al. Exosomes--Small Players, Big Sound. *Bioconjug Chem* 2018;29:635-48.
 46. Clayton A, Court J, Navabi H, et al. Analysis of antigen presenting cell derived exosomes, based on immunomagnetic isolation and flow cytometry. *J Immunol Methods* 2001;247:163-74.
 47. Carnell-Morris P, Tannetta D, Siupa A, et al. Analysis of Extracellular Vesicles Using Fluorescence Nanoparticle Tracking Analysis. *Methods Mol Biol* 2017;1660:153-73.
 48. Anderson W, Lane R, Korbie D, et al. Observations of Tunable Resistive Pulse Sensing for Exosome Analysis:

- Improving System Sensitivity and Stability. *Langmuir* 2015;31:6577-87.
49. Liu JJ, Bratkowski MA, Liu X, et al. Visualization of distinct substrate-recruitment pathways in the yeast exosome by EM. *Nat Struct Mol Biol* 2014;21:95-102.
 50. Keller S, Sanderson MP, Stoeck A, et al. Exosomes: from biogenesis and secretion to biological function. *Immunol Lett* 2006;107:102-8.
 51. Buschow SI, van Balkom BW, Aalberts M, et al. MHC class II-associated proteins in B-cell exosomes and potential functional implications for exosome biogenesis. *Immunol Cell Biol* 2010;88:851-6.
 52. Stoorvogel W, Kleijmeer MJ, Geuze HJ, et al. The biogenesis and functions of exosomes. *Traffic* 2002;3:321-30.
 53. Baietti MF, Zhang Z, Mortier E, et al. Syndecan-syntenin-ALIX regulates the biogenesis of exosomes. *Nat Cell Biol* 2012;14:677-85.
 54. Kahlert C, Kalluri R. Exosomes in tumor microenvironment influence cancer progression and metastasis. *J Mol Med (Berl)* 2013;91:431-7.
 55. Utsugi-Kobukai S, Fujimaki H, Hotta C, et al. MHC class I-mediated exogenous antigen presentation by exosomes secreted from immature and mature bone marrow derived dendritic cells. *Immunol Lett* 2003;89:125-31.
 56. Clayton A, Turkes A, Dewitt S, et al. Adhesion and signaling by B cell-derived exosomes: the role of integrins. *FASEB J* 2004;18:977-9.
 57. Rieu S, Géminard C, Rabesandratana H, et al. Exosomes released during reticulocyte maturation bind to fibronectin via integrin alpha4beta1. *Eur J Biochem* 2000;267:583-90.
 58. Hao S, Bai O, Li F, et al. Mature dendritic cells pulsed with exosomes stimulate efficient cytotoxic T-lymphocyte responses and antitumour immunity. *Immunology* 2007;120:90-102.
 59. Morelli AE, Larregina AT, Shufesky WJ, et al. Endocytosis, intracellular sorting, and processing of exosomes by dendritic cells. *Blood* 2004;104:3257-66.
 60. Fitzner D, Schnaars M, van Rossum D, et al. Selective transfer of exosomes from oligodendrocytes to microglia by macropinocytosis. *J Cell Sci* 2011;124:447-58.
 61. Feng D, Zhao WL, Ye YY, et al. Cellular internalization of exosomes occurs through phagocytosis. *Traffic* 2010;11:675-87.
 62. Nanbo A, Kawanishi E, Yoshida R, et al. Exosomes derived from Epstein-Barr virus-infected cells are internalized via caveola-dependent endocytosis and promote phenotypic modulation in target cells. *J Virol* 2013;87:10334-47.
 63. Svensson KJ, Christianson HC, Wittrup A, et al. Exosome uptake depends on ERK1/2-heat shock protein 27 signaling and lipid Raft-mediated endocytosis negatively regulated by caveolin-1. *J Biol Chem* 2013;288:17713-24.
 64. Escrevente C, Keller S, Altevogt P, et al. Interaction and uptake of exosomes by ovarian cancer cells. *BMC Cancer* 2011;11:108.
 65. Hoshino A, Costa-Silva B, Shen TL, et al. Tumour exosome integrins determine organotropic metastasis. *Nature* 2015;527:329-35.
 66. Zhang Y, Hu YW, Zheng L, et al. Characteristics and Roles of Exosomes in Cardiovascular Disease. *DNA Cell Biol* 2017;36:202-11.
 67. Jansen F, Li Q. Exosomes as Diagnostic Biomarkers in Cardiovascular Diseases. *Adv Exp Med Biol* 2017;998:61-70.
 68. Kanninen KM, Bister N, Koistinaho J, et al. Exosomes as new diagnostic tools in CNS diseases. *Biochim Biophys Acta* 2016;1862:403-10.
 69. Fitts CA, Ji N, Li Y, et al. Exploiting Exosomes in Cancer Liquid Biopsies and Drug Delivery. *Adv Healthc Mater* 2019;8:e1801268.
 70. Masyuk AI, Masyuk TV, Larusso NF. Exosomes in the pathogenesis, diagnostics and therapeutics of liver diseases. *J Hepatol* 2013;59:621-5.
 71. Zhang W, Zhou X, Zhang H, et al. Extracellular vesicles in diagnosis and therapy of kidney diseases. *Am J Physiol Renal Physiol* 2016;311:F844-F851.
 72. Alipoor SD, Mortaz E, Varahram M, et al. The Potential Biomarkers and Immunological Effects of Tumor-Derived Exosomes in Lung Cancer. *Front Immunol* 2018;9:819.
 73. Zhou X, Xie F, Wang L, et al. The function and clinical application of extracellular vesicles in innate immune regulation. *Cell Mol Immunol* 2020;17:323-34.
 74. Cai J, Han Y, Ren H, et al. Extracellular vesicle-mediated transfer of donor genomic DNA to recipient cells is a novel mechanism for genetic influence between cells. *J Mol Cell Biol* 2013;5:227-38.
 75. Kahlert C, Melo SA, Protopopov A, et al. Identification of double-stranded genomic DNA spanning all chromosomes with mutated KRAS and p53 DNA in the serum exosomes of patients with pancreatic cancer. *J Biol Chem* 2014;289:3869-75.
 76. Kalluri R, LeBleu VS. Discovery of Double-Stranded Genomic DNA in Circulating Exosomes. *Cold Spring Harb Symp Quant Biol* 2016;81:275-80.
 77. Thakur BK, Zhang H, Becker A, et al. Double-stranded DNA in exosomes: a novel biomarker in cancer detection.

- Cell Res 2014;24:766-9.
78. Salehi M, Sharifi M. Exosomal miRNAs as novel cancer biomarkers: Challenges and opportunities. *J Cell Physiol* 2018;233:6370-80.
 79. Thind A, Wilson C. Exosomal miRNAs as cancer biomarkers and therapeutic targets. *J Extracell Vesicles* 2016;5:31292.
 80. Lian J, Lin SH, Ye Y, et al. Serum microRNAs as predictors of risk for non-muscle invasive bladder cancer. *Oncotarget* 2018;9:14895-908.
 81. Hallal S, Azimi A, Wei H, et al. A Comprehensive Proteomic SWATH-MS Workflow for Profiling Blood Extracellular Vesicles: A New Avenue for Glioma Tumour Surveillance. *Int J Mol Sci* 2020;21:4754.
 82. Halvaei S, Daryani S, Eslami-S Z, et al. Exosomes in Cancer Liquid Biopsy: A Focus on Breast Cancer. *Mol Ther Nucleic Acids* 2018;10:131-41.
 83. Stevic I, Müller V, Weber K, et al. Specific microRNA signatures in exosomes of triple-negative and HER2-positive breast cancer patients undergoing neoadjuvant therapy within the GeparSixto trial. *BMC Med* 2018;16:179.
 84. Zhou X, Zhu W, Li H, et al. Diagnostic value of a plasma microRNA signature in gastric cancer: a microRNA expression analysis. *Sci Rep* 2015;5:11251.
 85. Yosef I, Goren MG, Qimron U. Proteins and DNA elements essential for the CRISPR adaptation process in *Escherichia coli*. *Nucleic Acids Res* 2012;40:5569-76.
 86. Cho E, Nam GH, Hong Y, et al. Comparison of exosomes and ferritin protein nanocages for the delivery of membrane protein therapeutics. *J Control Release* 2018;279:326-35.
 87. György B, Hung ME, Breakefield XO, et al. Therapeutic applications of extracellular vesicles: clinical promise and open questions. *Annu Rev Pharmacol Toxicol* 2015;55:439-64.
 88. Tian T, Zhang HX, He CP, et al. Surface functionalized exosomes as targeted drug delivery vehicles for cerebral ischemia therapy. *Biomaterials* 2018;150:137-49.
 89. Rani S, Ryan AE, Griffin MD, et al. Mesenchymal Stem Cell-derived Extracellular Vesicles: Toward Cell-free Therapeutic Applications. *Mol Ther* 2015;23:812-23.
 90. Zhang B, Yin Y, Lai RC, et al. Mesenchymal stem cells secrete immunologically active exosomes. *Stem Cells Dev* 2014;23:1233-44.
 91. Lee C, Mitsialis SA, Aslam M, et al. Exosomes mediate the cytoprotective action of mesenchymal stromal cells on hypoxia-induced pulmonary hypertension. *Circulation* 2012;126:2601-11.
 92. Kanazawa H, Fujimoto Y, Teratani T, et al. Bone Marrow-Derived Mesenchymal Stem Cells Ameliorate Hepatic Ischemia Reperfusion Injury in a Rat Model. *PLoS One* 2011;6:e19195.
 93. Lai RC, Arslan F, Lee MM, et al. Exosome secreted by MSC reduces myocardial ischemia/reperfusion injury. *Stem Cell Res* 2010;4:214-22.
 94. Salomon C, Ryan J, Sobrevia L, et al. Exosomal Signaling during Hypoxia Mediates Microvascular Endothelial Cell Migration and Vasculogenesis. *PLoS One* 2013;8:e68451.
 95. Chen J, Liu Z, Hong MM, et al. Proangiogenic compositions of microvesicles derived from human umbilical cord mesenchymal stem cells. *PLoS One* 2014;9:e115316.
 96. Yang Y, Hong Y, Cho E, et al. Extracellular vesicles as a platform for membrane-associated therapeutic protein delivery. *J Extracell Vesicles* 2018;7:1440131.
 97. Bunggulawa EJ, Wang W, Yin T, et al. Recent advancements in the use of exosomes as drug delivery systems. *J Nanobiotechnology* 2018;16:81.
 98. Jan AT, Malik MA, Rahman S, et al. Perspective Insights of Exosomes in Neurodegenerative Diseases: A Critical Appraisal. *Front Aging Neurosci* 2017;9:317.
 99. ClinicalTrials.gov [Accessed on 23.07.2020]. Available online: https://clinicaltrials.gov/ct2/results?cond=exosome&Search=Apply&recrs=e&age_v=&gndr=&type=&rslt=
 100. Wang K, Jiang Z, Webster KA, et al. Enhanced Cardioprotection by Human Endometrium Mesenchymal Stem Cells Driven by Exosomal MicroRNA-21. *Stem Cells Transl Med* 2017;6:209-22.
 101. Lou G, Chen Z, Zheng M, et al. Mesenchymal stem cell-derived exosomes as a new therapeutic strategy for liver diseases. *Exp Mol Med* 2017;49:e346.
 102. Kim HS, Choi DY, Yun SJ, et al. Proteomic analysis of microvesicles derived from human mesenchymal stem cells. *J Proteome Res* 2012;11:839-49.
 103. Merino-González C, Zuñiga FA, Escudero C, et al. Mesenchymal Stem Cell-Derived Extracellular Vesicles Promote Angiogenesis: Potential Clinical Application. *Front Physiol* 2016;7:24.
 104. Tracy SA, Ahmed A, Tigges JC, et al. A comparison of clinically relevant sources of mesenchymal stem cell-derived exosomes: Bone marrow and amniotic fluid. *J Pediatr Surg* 2019;54:86-90.
 105. Ha D, Yang N, Nadithe V. Exosomes as therapeutic drug

- carriers and delivery vehicles across biological membranes: current perspectives and future challenges. *Acta Pharm Sin B* 2016;6:287-96.
106. Di Rocco G, Baldari S, Toietta G. Towards Therapeutic Delivery of Extracellular Vesicles: Strategies for In Vivo Tracking and Biodistribution Analysis. *Stem Cells Int* 2016;2016:5029619.
 107. Ren S, Chen J, Duscher D, et al. Microvesicles from human adipose stem cells promote wound healing by optimizing cellular functions via AKT and ERK signaling pathways. *Stem Cell Res Ther* 2019;10:47.
 108. Zhang K, Yu L, Li FR, et al. Topical Application of Exosomes Derived from Human Umbilical Cord Mesenchymal Stem Cells in Combination with Sponge Spicules for Treatment of Photoaging. *Int J Nanomedicine* 2020;15:2859-72.
 109. Ferreira ADF, Gomes DA. Stem Cell Extracellular Vesicles in Skin Repair. *Bioengineering (Basel)* 2018;6:4.
 110. Sengupta V, Sengupta S, Lazo A, et al. Exosomes Derived from Bone Marrow Mesenchymal Stem Cells as Treatment for Severe COVID-19. *Stem Cells Dev* 2020;29:747-54.
 111. Ilic N, Savic S, Siegel E, et al. Examination of the regulatory frameworks applicable to biologic drugs (including stem cells and their progeny) in Europe, the U.S., and Australia: Part II—A method of software documentary analysis. *Stem Cells Transl Med* 2012;1:909-20.
 112. Ilic N, Savic S, Siegel E, et al. Examination of the regulatory frameworks applicable to biologic drugs (including stem cells and their progeny) in Europe, the U.S., and Australia: part I—a method of manual documentary analysis. *Stem Cells Transl Med* 2012;1:898-908.
 113. European_Union. Volume 4 EU Guidelines for Good Manufacturing Practice for Medicinal Products for Human and Veterinary Use; Annex 2, Manufacture of Biological Active Substances and Medicinal Products for Human Use.
 114. GMPguidelines. [assessed on 28 July 2020]. Available online: http://ec.europa.eu/health/documents/eudralex/vol-4/index_en.htm
 115. ClinicalTrialsGuidelines. [assessed on 28 July 2020]. Available online: http://ec.europa.eu/health/documents/eudralex/vol-10/index_en.htm
 116. EMA/CHMP/BWP/534898/2008. Guideline on the requirements for quality documentation concerning biological investigational medicinal products in clinical trials.
 117. Lener T, Gimona M, Aigner L, et al. Applying extracellular vesicles based therapeutics in clinical trials - an ISEV position paper. *J Extracell Vesicles* 2015;4:30087.
 118. EMEA/CHMP/410869/2006. Guideline on human cell-based medicinal products.
 119. Ruiss R, Jochum S, Mocikat R, et al. EBV-gp350 Confers B-Cell Tropism to Tailored Exosomes and Is a Neo-Antigen in Normal and Malignant B Cells—A New Option for the Treatment of B-CLL. *PLoS One* 2011;6:e25294.
 120. REGULATION/EC/1394/2007. Regulation on advanced therapy medicinal products.
 121. EMA/CAT/CPWP/686637/2011. Guideline on the risk-based approach according to annex I, part IV of Directive 2001/83/EC applied to Advanced therapy medicinal products.
 122. Kriebardis A, Antonelou M, Stamoulis K, et al. Cell-derived microparticles in stored blood products: innocent-bystanders or effective mediators of post-transfusion reactions? *Blood Transfus* 2012;10 Suppl 2:s25-s38.
 123. Burke J, Kolhe R, Hunter M, et al. Stem Cell-Derived Exosomes: A Potential Alternative Therapeutic Agent in Orthopaedics. *Stem Cells Int* 2016;2016:5802529.
 124. Zhang J, Guan J, Niu X, et al. Exosomes released from human induced pluripotent stem cells-derived MSCs facilitate cutaneous wound healing by promoting collagen synthesis and angiogenesis. *J Transl Med* 2015;13:49.
 125. Devhare PB, Ray RB. A novel role of exosomes in the vaccination approach. *Ann Transl Med* 2017;5:23.
 126. Nolte-'t Hoen E, Cremer T, Gallo RC, et al. Extracellular vesicles and viruses: Are they close relatives? *Proc Natl Acad Sci U S A* 2016;113:9155-61.
 127. Joo HS, Suh JH, Lee HJ, et al. Current Knowledge and Future Perspectives on Mesenchymal Stem Cell-Derived Exosomes as a New Therapeutic Agent. *Int J Mol Sci* 2020;21:727.
 128. Capricor Therapeutics. [assessed on 28 July 2020]. Available online: <http://capricor.com/>
 129. Kimera Labs. [assessed on 28 July 2020]. Available online: <https://kimeralabs.com/>
 130. Exosome Diagnostics. Personalized Precision Healthcare. [assessed on 28 July 2020]. Available online: <http://www.exosomedx.com/>
 131. NanoSomiX Inc. [assessed on 28 July 2020]. Available online: <http://www.nanosomix.com/technology>
 132. Evox Therapeutics. [assessed on 28 July 2020]. Available online: <https://www.evoxtherapeutics.com/>
 133. Codiak Biosciences. Realizing the potential of exosome biology. [assessed on 28 July 2020]. Available online:

<http://www.codiakbio.com/#cellular-communicators>
134. ArunA Biomedical. [assessed on 28 July 2020]. Available
online: <https://www.arunabiomedical.com/>

135. de Couto G, Liu W, Tseliou E, et al. Macrophages
mediate cardioprotective cellular postconditioning in acute
myocardial infarction. *J Clin Invest* 2015;125:3147-62.

doi: 10.21037/sci-2020-037

Cite this article as: Muthu S, Bapat A, Jain R, Jeyaraman N,
Jeyaraman M. Exosomal therapy—a new frontier in
regenerative medicine. *Stem Cell Investig* 2021;8:7.